

Lil
UNCLASSIFIED

MA 2373
AD

2373
D 657 316

BODY VOLUME OF ADULT MEN

Chester L. Ward

**USAF School of Aerospace Medicine
Brooks Air Force Base, Texas**

June 1967

Processed for . . .

**DEFENSE DOCUMENTATION CENTER
DEFENSE SUPPLY AGENCY**



20030113002

U. S. DEPARTMENT OF COMMERCE / NATIONAL BUREAU OF STANDARDS / INSTITUTE FOR APPLIED TECHNOLOGY

UNCLASSIFIED

NOTICE TO DEFENSE DOCUMENTATION CENTER USERS

This document is being distributed by the Clearinghouse for Federal Scientific and Technical Information, Department of Commerce, as a result of a recent agreement between the Department of Defense (DOD) and the Department of Commerce (DOC).

The Clearinghouse is distributing unclassified, unlimited documents which are or have been announced in the Technical Abstract Bulletin (TAB) of the Defense Documentation Center.


The price does not apply for registered users of the DDC services.

AD657316

BODY VOLUME OF ADULT MEN

CHESTER L. WARD, Major, MC, F/S, USA

June 1967



USAF School of Aerospace Medicine
Aerospace Medical Division (AFSC)
Brooks Air Force Base, Texas

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

BODY VOLUME OF ADULT MEN

CHESTER L. WARD, Major, MC, F/S, USA


This document has been approved
for public release and sale; its
distribution is unlimited.

FOREWORD

The research reported in this paper was conducted by personnel at the USAF School of Aerospace Medicine, Brooks AFB, Tex., under task No. 775801. The measurements for this report were obtained from 10 March 1964 through 9 December 1966 by personnel in the Physiology Branch. The report was submitted for publication on 6 March 1967.

The author gratefully acknowledges the assistance of the following persons in the Physiology and Biometrics Branches of the School: Dr. Thomas H. Allen, for technical advice and consultation; Clarence Theis, for acquisition of measurements; and Dr. Bryan Danford and CMSgt. James I. Clogston for data processing, analysis, and interpretation.

This report has been reviewed and is approved.


JAMES E. NUTTALL
Colonel, USAF, MC
Commander

ABSTRACT

Body composition determinations were made on 404 adult men by use of a volumetric method. The testing of a proposed nomogram for estimation of body volume from height and weight revealed the chart to be biased for adult men. Body volume was found to correlate well with body weight (correlation coefficient of .996). Body volume of men in liters, V , may be estimated from body weight in kilograms, W , by using the formula: $V = -4.7573 + 1.0153 W$.

The ideal weight given on the USAF standard weight table was found to have a correlation coefficient of only .672 with calculated percent body fat.

BODY VOLUME OF ADULT MEN

I. INTRODUCTION

Human body composition is a subject of great interest and continuing research. The component of greatest interest and that which appears to be of the greatest importance relative to health, morbidity, and mortality is the adipose tissue (8). It is often important to the armed forces, insurance companies, and other industries—as well as to the person concerned about his own appearance, efficiency, and health—to distinguish between having excessive adipose tissue and merely being “overweight.” An individual may be overweight according to standard height-weight tables and still not be obese (11). This relatively increased weight may be the result of an athletic, heavy, muscular physique rather than obesity or “overfat.”

The purpose of this report is to present the anthropometric data of 404 adult men, aged 17 to 51 years. These body measurements and computations of body composition will be compared and contrasted with the USAF weight table (1) and with a method recently suggested by Sendroy and Collison (10), using a nomogram, for predicting body volume from measurements of height and weight. A formula for estimating body volume from weight alone will be presented.

II. METHOD

Measurements were made and recorded for each subject's age, height, weight, observed body volume, and predicted body volume. On the basis of these measurements, the following calculations were made for the data on each subject: density, difference between measured and predicted volume, fat, percent of fat to weight, and percent of relative ideal weight.

Certain descriptive statistics were derived from these data for presentation with the results.

Body composition was determined by a volumetric method (2) with appropriate deduction of measured residual lung volume. This method assumes that the greatest variable in adult human body composition is adipose tissue and that the densities of body components are relatively constant. Discussion of the validity of these premises exceeds the purpose of this report; reviews on body composition, in which these ideas are discussed, are available in the literature (4-7, 9).

Subjects

U. S. Air Force personnel on active duty constituted most of the 404 adult males measured. Significant groups of subjects included the following: normal men exposed to experimental aerospace atmospheres, altitudes, confined quarters, and foods—142; obese subjects for special diets—64; MOL astronaut candidates—45; physicians in Aerospace Medicine Residency—37; patients seen in consultation having diseases other than obesity—23; U.S. pentathlon team—14; and “skinny” men, subjects for a high fat diet study—8. The only observations used in this report are the measurements recorded the first time a subject presented himself, in his own control state, for study; repeat values and subsequent changes in an individual are excluded from this report.

Age. The subject gave his age as years at last anniversary of birth.

Height. Measurement was obtained to the nearest millimeter as the subject stood barefooted with heels together and stretched to maximum height.

Weight. The nude subject, fasting or after a very light breakfast, was weighed dry in the morning after a shower, micturition, and defecation, if possible. Weight was determined to the nearest 5 gm. on a standard beam balance scale.

Body volume. The subject's volume was measured by water displacement using the technic and volumeter developed by Allen (2). After the subject was weighed, as described above, and before he was immersed in the volumeter, his residual lung volume was determined to the nearest mean cubic millimeter on three consistent trials by the method using open circuit nitrogen washout (with oxygen). Body volume of tissue was determined to the nearest cubic millimeter by subtracting the subject's measured residual lung volume from his submerged volume after complete underwater expiration of his vital capacity.

Predicted body volume. A nomogram proposed by Sendroy and Collison (10) for determining body volume was tested. Predicted volume was determined by interpolation to the nearest liter from an enlarged nomographic chart in the manner recommended by the authors named above. Predicted volumes were determined by an independent second reading and all differences in interpolation were arbitrated.

Difference. Measured volume minus the predicted volume ($\Delta = V_o - V_p$) was calculated.

Density. Weight divided by measured volume was calculated to six decimal places ($D = W/V$).

Percent fat. Weight of fat was calculated using the formula $F = c_1V - c_2W$ where F is fat in kilograms, $c_1 = 4.878$ (kg. liter⁻¹), V is volume in liters, $c_2 = 4.415$ (kg. kg.⁻¹), and W is weight in kilograms (3). Weight of the body fat was divided by total body weight and multiplied by 100 ($\%F = 100 [F/W]$).

Percent ideal weight. The "ideal weight" used was the standard weight for the 26- to 30-year age group in the USAF weight table (1). The difference of observed and ideal weight divided by ideal weight, all multiplied by 100, is defined as percent ideal weight ($\%Wi = 100 [(W_o - W_i)/W_i]$).

III. RESULTS

Descriptive data on the 404 subjects are presented in table I and data means for 5-year age groupings are presented in table II.

The expected increase in the mean weight with advancing age is noted in the 5-year age

TABLE I
Descriptive data, 404 subjects

Variable	Mean	Range		S. D.
		Minimum	Maximum	
Age, years*	31.70	17	51	8.06
Height, cm.	178.10	157.7	196	5.86
Weight, kg.	77.61	47.59	117.98	10.95
Volume, liters	74.05	44.31	119.15	11.16
Density	1.050	.990	1.097	.016
Percent fat	23.31	3.21	51.13	7.06
Percent ideal weight	9.68	-24.52	72.25	13.38

*Number for age grouping in 404: one age was not recorded.

groupings except for the 41- to 45-year aged subject grouping. The reason for the deviation from the progression of increasing weight, volume, percent fat, and percent ideal weight and the decreasing density observed in the means for the 41- to 45-year age group is unknown.

The testing of a chart-nomogram method (10) for predicting body volume is presented in the "Liters difference" row of table II. The enlarged nomographic chart was found relatively difficult to read and, at best, can be interpolated only to the nearest full liter of predicted body volume. The mean difference between the measured body volume and predicted volume for the 404 subjects was 1.755 liters with a standard deviation of 1.234 liters.

If a *rough* body volume estimate is needed for such purposes as calculating predicted body fat (see formula above) or for other clinical and practical purposes, body volume may be estimated for adult men using the formula, $V = -4.757 + 1.015 W$, derived from the plot of the regression illustrated by the scattergram in figure 1. The correlation coefficient of weight and volume is .996; thus, volume in liters, V, may be fairly accurately predicted from weight in kilograms, W. If one uses the formula $V = -4.757 + 1.015 W$ to predict

volume from weight, then the 95% confidence limits on the predicted volume, V_p , is $V_p \pm 1.9$ liters.

Percent ideal weight has correlation coefficients with density and percent fat of only $-.672$ and $+.672$, respectively. It should be observed that the information content of density and percent fat is the same. This relationship may be verified algebraically as well as empirically, as shown in the correlation coefficients above.

IV. DISCUSSION

The preceding results have demonstrated that a recently proposed nomogram for the determination of human body volume from height and weight is biased. A more accurate and easier-to-use formula for estimating the body volume of adult males has been presented.

The estimation of fatness in the 404 measured subjects by equating fatness or obesity with the concept of overweight, as compared to an "ideal" weight from a height-weight table, revealed a correlation coefficient of only .67 between calculated percent fat and departure from ideal weight. This relatively poor correlation between measured fatness and a weight table indicates that charts or tables based upon

TABLE II
Means of variables in 5-year age groups

Variable	16-20	21-25	26-30	31-35	36-40	41-45	46-50	51-55
Height, cm.	179.0	178.3	178.4	177.9	177.4	177.1	179.4	182.5
Weight, kg.	72.30	75.39	78.69	78.90	79.42	75.86	83.04	91.17
Volume, liters	68.10	71.41	74.96	75.31	76.34	72.75	80.07	87.94
Liters predicted	67.20	70.30	73.24	73.53	73.87	70.58	77.88	86.00
Liters difference	.90	1.11	1.72	1.78	2.47	2.17	2.19	1.94
Density	1.064	1.058	1.051	1.049	1.041	1.044	1.038	1.037
Percent fat	17.06	19.74	22.73	23.72	27.05	25.99	28.52	29.00
Percent ideal weight	1.56	6.20	10.78	11.66	13.15	8.42	15.58	22.08
Number subjects*	35	63	94	81	52	59	17	2

*See footnote, table I.




FIGURE 1

Scatter diagram showing relationship between volume and weight.

age, height, and weight are an inadequate method for the evaluation of a person's body composition. Overweight means only that when compared to an observed mean weight from a population with the subject's same height and

age, the subject has somewhat more than the expected weight. The subject's relatively high weight might very well be due to a heavy skeletal system and a large, well-developed, muscle mass rather than excess fat or obesity as the term *overweight* implies—often incorrectly.

The concept that obesity rather than overweight may adversely affect health and longevity requires that methods better than reference to age-height-weight tables be used to estimate body composition. Clinical judgment, testing of functional capacity, and measurement of skin-fold thickness are quite useful; but for exceptional individuals (those who do not conform to the model or physical stereotype) additional and more accurate methods of determining body composition are necessary.

The volumetric method for determining density and estimating body fat, using a standard table to estimate residual lung volume if necessary, is readily usable in nearly any situation because it requires no complex equipment, is relatively simple to operate, and has a low initial cost with minimal upkeep. A volumeter could be easily fabricated from materials available in most remote or undeveloped areas. The use of this volumeter would give an accurate method of determining body composition.

REFERENCES

1. Air Force Manual 160-1. Medical examination and medical standards. Attachment 1, p. 111. Washington, D.C.: U.S. Government Printing Office, 1964.
2. Allen, T. H. Measurement of human body fat: A quantitative method suited for use by aviation medical officers. *Aerospace Med.* 34:907-909 (1963).
3. Allen, T. H. Personal communication, 5 August 1966.
4. Brozek, J. Human body composition: Approaches and applications. (Symposia of the Society for the Study of Human Biology, vol. 7) New York: Pergamon Press, 1965.
5. Brozek, J. Body composition conference by the New York Academy of Sciences. *Ann. N.Y. Acad. Sci.*, vol. 110 (1963).
6. Brozek, J., and A. Henschel. Conference on Techniques for Measuring Body Composition, Natick, Mass., 1959. Washington, D.C.: National Academy of Sciences—National Research Council, 1961.
7. Forbes, G. B. Methods for determining composition of the human body. *Pediatrics* 29:477-494 (1962).
8. Fox, S. M., III. Obesity and health, a source book of current information for professional health personnel. Public Health Service Publication No. 1485, Washington, D.C.: U.S. Government Printing Office, 1966.
9. Keys, A., and J. Brozek. Body fat in adult man. *Physiol. Rev.* 33:245-325 (1953).
10. Sendroy, J., Jr., and H. A. Collison. Determination of human body volume from height and weight. *J. Appl. Physiol.* 21:167-172 (1966).
11. Welham, W. C., and A. R. Behnke, Jr. The specific gravity of healthy men. Body weight ÷ volume and other physical characteristics of exceptional athletes and of naval personnel. *J.A.M.A.* 118:498-501 (1942).

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) USAF School of Aerospace Medicine Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP
3. REPORT TITLE BODY VOLUME OF ADULT MEN		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) 10 Mar. 1964 - 9 Dec. 1966		
5. AUTHOR(S) (Last name, first name, initial) Ward, Chester L., Major, MC, USA		
6. REPORT DATE June 1967	7a. TOTAL NO. OF PAGES 5	7b. NO. OF REFS 11
8a. CONTRACT OR GRANT NO.	9a. ORIGINATOR'S REPORT NUMBER(S) SAM-TR-67-42	
b. PROJECT NO.		
c. Task No. 775801	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10. AVAILABILITY/LIMITATION NOTICES This document has been approved for public release and sale; its distribution is unlimited.		
11. SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY USAF School of Aerospace Medicine Aerospace Medical Division (AFSC) Brooks Air Force Base, Texas	
13. ABSTRACT Body composition determinations were made on 404 adult men by use of a volumetric method. The testing of a proposed nomogram for estimation of body volume from height and weight revealed the chart to be biased for adult men. Body volume was found to correlate well with body weight (correlation coefficient of .996). Body volume of men in liters, V, may be estimated from body weight in kilograms, W, by using the formula: $V = -4.7573 + 1.0153 W$. The ideal weight given on the USAF standard weight table was found to have a correlation coefficient of only .672 with calculated percent body fat.		

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Body volume (volumetric method) Nomogram, body volume Weight tables Obesity vs. overweight						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.